



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2020

---

## **Detection of Chlamydiaceae in Swiss wild birds sampled at a bird rehabilitation centre**

Stalder, Sandro ; Marti, Hanna ; Borel, Nicole ; Mattmann, Prisca ; Vogler, Barbara Renate ; Wolfrum, Nina ; Albin, Sarah

**Abstract:** Background: Annually, 800-1500 wild birds are admitted to the rehabilitation centre of the Swiss Ornithological Institute, Sempach, Lucerne, Switzerland. The workers of the centre come in close contact with the avian patients and might therefore be exposed to zoonotic agents shed by these birds, such as *Chlamydia psittaci*. **Methods:** In the present study, 91 choanal, 91 cloacal and 267 faecal swabs from 339 wild birds of 42 species were investigated using a stepwise diagnostic approach. **Results:** Chlamydiaceae were detected in 0.9 per cent (0.3-2.6 per cent) of birds (n=3), all of them members of the Columbidae family. The Chlamydiaceae species of two of these birds (one Eurasian collared dove, one fancy pigeon) were identified as *C. psittaci* types B and E by PCR and outer membrane protein A genotyping. **Conclusion:** The findings of the current study suggest that zoonotic transmission of Chlamydiaceae is very unlikely for songbird and waterfowl species tested herein, while pigeons might pose a risk to workers at rehabilitation centres.

DOI: <https://doi.org/10.1136/vetreco-2020-000437>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-195722>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

Originally published at:

Stalder, Sandro; Marti, Hanna; Borel, Nicole; Mattmann, Prisca; Vogler, Barbara Renate; Wolfrum, Nina; Albin, Sarah (2020). Detection of Chlamydiaceae in Swiss wild birds sampled at a bird rehabilitation centre. *Veterinary record open*, 7(1):e000437.

DOI: <https://doi.org/10.1136/vetreco-2020-000437>

# Detection of *Chlamydiaceae* in Swiss wild birds sampled at a bird rehabilitation centre

Sandro Stalder ,<sup>1</sup> Hanna Marti,<sup>2</sup> Nicole Borel,<sup>2</sup> Prisca Mattmann,<sup>3</sup> Barbara Vogler ,<sup>1</sup> Nina Wolfrum,<sup>1</sup> Sarah Albini<sup>1</sup>

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/vetreco-2020-000437>).

**To cite:** Stalder S, Marti H, Borel N, *et al.* Detection of *Chlamydiaceae* in Swiss wild birds sampled at a bird rehabilitation centre. *Veterinary Record Open* 2020;**7**:e000437. doi:10.1136/vetreco-2020-000437

Received 27 July 2020  
Revised 23 September 2020  
Accepted 13 October 2020

## ABSTRACT

**Background** Annually, 800–1500 wild birds are admitted to the rehabilitation centre of the Swiss Ornithological Institute, Sempach, Lucerne, Switzerland. The workers of the centre come in close contact with the avian patients and might therefore be exposed to zoonotic agents shed by these birds, such as *Chlamydia psittaci*.

**Methods** In the present study, 91 cloacal, 91 cloacal and 267 faecal swabs from 339 wild birds of 42 species were investigated using a stepwise diagnostic approach.

**Results** *Chlamydiaceae* were detected in 0.9 per cent (0.3–2.6 per cent) of birds (n=3), all of them members of the Columbidae family. The *Chlamydiaceae* species of two of these birds (one Eurasian collared dove, one fancy pigeon) were identified as *C psittaci* types B and E by PCR and outer membrane protein A genotyping.

**Conclusion** The findings of the current study suggest that zoonotic transmission of *Chlamydiaceae* is very unlikely for songbird and waterfowl species tested herein, while pigeons might pose a risk to workers at rehabilitation centres.

## INTRODUCTION

The rehabilitation centre of the Swiss Ornithological Institute is located in Sempach, Lucerne, Switzerland. In the last 10 years, between 800 and 1500 birds were admitted for treatment annually. The workers of the rehabilitation centre come in close contact with the birds and their faeces during the time of treatment. Thus, the workers might be exposed to zoonotic agents shed by these birds, such as *Chlamydia psittaci*. Several *Chlamydiaceae* species are harboured by wild birds, for example, *C abortus*, *C avium*, *C pecorum* and *Candidatus C ibidis*.<sup>1–7</sup> *C psittaci*, the causative agent of psittacosis/ornithosis in human and avian chlamydiosis, has been detected in more than 460 avian species.<sup>8</sup> The clinical signs in *C psittaci*-infected birds are variable, depending on age, species and immune status of the host, and the pathogenicity of the causative strain.<sup>9</sup> Clinical signs include respiratory, enteric and ocular signs, but asymptomatic infections are common.<sup>10</sup> In Switzerland, 46 cases in birds

have been reported to the Federal Food Safety and Veterinary Office from 2010 to 2019.<sup>11</sup> Transmission from birds to humans occurs through inhalation of feather dust or aerosols from urine, dried faeces, and respiratory and eye secretions.<sup>12</sup> Most infected humans remain asymptomatic or have mild symptoms, but in some cases *C psittaci* causes severe pneumonia.

Studies investigating the occurrence of *Chlamydiaceae* in European wild birds reported infection rates ranging from 2.8 per cent to 14.8 per cent determined using PCR or immunoassay.<sup>5 13–18</sup> In Switzerland, two PCR-based studies focusing on pigeons reported infection rates of 8.4 per cent and 16.9 per cent,<sup>19 20</sup> and another PCR-based study reported infection rates of 14.3 per cent in pigeons, 0.4 per cent in songbirds and 4.3 per cent in waterfowl.<sup>21</sup> The highest infection rates of *Chlamydiaceae* were observed in Columbiformes and Psittaciformes, where *C psittaci* is the most frequently detected species.<sup>22 23</sup> Passeriformes were only occasionally diagnosed with chlamydiosis.<sup>24 25</sup>

The current study aims to investigate whether admitted wild birds pose a risk of zoonotic chlamydial infection to workers at a Swiss wild bird rehabilitation centre.

## MATERIALS AND METHODS

### Samples

Samples were obtained from the wild bird rehabilitation centre of the Swiss Ornithological Institute in Sempach, Lucerne, Switzerland, between April and October 2018 (online supplemental table 1). Around two-thirds of the birds were admitted during the peak season from May to mid-July, the majority of which were juveniles. Submitted avian patients received care depending on their condition, for example, feeding, cleaning, veterinary care, treatment or surgery if necessary. The staff of the rehabilitation centre



© British Veterinary Association 2020. Re-use permitted under CC BY-NC. No commercial re-use. Published by BMJ.

<sup>1</sup>National Reference Centre for Poultry and Rabbit Diseases, Institute for Food Safety and Hygiene, University of Zurich, Zurich, Switzerland

<sup>2</sup>Institute of Veterinary Pathology, University of Zurich, Zurich, Switzerland

<sup>3</sup>Swiss Ornithological Institute, Sempach, Lucerne, Switzerland

### Correspondence to

Sandro Stalder; [sandro.stalder07@gmail.com](mailto:sandro.stalder07@gmail.com)

(six persons, among one veterinarian) come into close contact with the birds while handling them, although manual handling is reduced to the necessary minimum. Juvenile birds are kept in small boxes with household paper or foam pads as substrate. Fledged birds are transferred to cages with foam pads or an outdoor aviary. The boxes and foam pads are cleaned every one to two hours using an industrial dishwasher; cages are cleaned with water and soap—depending on the amount and quality of faeces—daily or twice daily. Disinfection of the cages (Meliseptol rapid, B Braun) is carried out before each reassignment. The outdoor aviaries are cleaned when necessary using a high-pressure cleaner at 60°C while wearing a protective mask and goggles.

This study included 339 wild birds representing nine orders and 42 species, as shown in table 1.

The age of 316 birds was known, of which 163 were nestlings, 126 juveniles and 27 adults. Dry choanal (n=91) and cloacal swabs (n=91) (FLOQSwabs, COPAN Flock Technologies) were obtained from deceased birds only. Dry faecal swabs (n=267) were taken from living birds after defecation. Samples were stored at -80°C until further processing.

### DNA extraction

A commercial kit (Genomic DNA from tissue, NucleoSpin Tissue from Macherey-Nagel) was used to extract the DNA of choanal and cloacal swabs. DNA of the faecal samples was extracted with the Macherey-Nagel NucleoSpin stool kit according to the manufacturer's instructions. Extracted DNA was stored at -20°C until further use.

### Real-time qPCR assays for detection of *Chlamydiaceae*

First, all samples were tested in duplicates with a 23S rRNA-based *Chlamydiaceae* family-specific quantitative PCR (qPCR) (111 bp) modified to include an internal positive amplification control (IPC; enhanced green fluorescent protein (eGFP))<sup>26–28</sup> on an Applied Biosystems 7500 Real-Time PCR System (Thermo Fisher Scientific). The cycle conditions were 95°C for 20 seconds, followed by 45 cycles of 95°C for three seconds and 60°C for 30 seconds. For every sample, a 25 µl reaction mix was prepared, including 12.5 µl TaqMan Universal PCR MasterMix, 500 nM of the primers 'CH23S-F' (5'-CTGA AACCAGTAGCTTATAAGCGGT-3') and 'CH23S-R' (5'-ACCTCGCCGTTTAACTTAACTCC-3'), 200 nM of the probe 'CH23S-P' (5'-FAM-CTCATCATGCAAAAGG CACGCCG-TAMRA 3'), and 200 nM each of the primers 'eGFP-1-F' (5'-GACCACTACCAGCAGAACAC-3') and 'eGFP-10-R' (5'-CTTGATACAGCTCGTCCATGC-3'), and the probe 'eGFP-HEX' (5'-HEX-AGCACCCAGTCC GCCCTGAGCA-BHQ1-3'). A sevenfold dilution series of *C. abortus* DNA with a known number of DNA copies was included in each run as a positive control and standard curve. Molecular grade water was included as a negative control in each run. Samples were interpreted as positive if the mean cycle threshold (Ct value) was less than 38.

Samples with higher Ct values or inhibited amplification were retested in duplicate. Samples repeatedly showing a Ct value greater than 38 were considered positive. Samples with inhibited amplification were retested undiluted and tenfold diluted, both in duplicates.

Secondly, in samples positive for *Chlamydiaceae*, a *C. psittaci*-specific qPCR (76 bp) was performed as previously described, modified to include an IPC.<sup>29,30</sup> The reaction mix contained 4 µl (<150 ng/µl) sample template, 1 µl eGFP template, 1x TaqMan Universal PCR MasterMix, 900 nM of the primers 'CppsOMP1-F' (5'-CACTATGT GGAAGGTGCTTCA-3') and 'CppsOMP1-R' (5'-CTGC-GCGGATGCTAATGG-3'), 200 nM probe 'CppsOMP1-S' (5'-FAM-CGCTACTTGGTGTGAC-TAMRA-3'), 900 nM of the primers 'eGFP-1-F' (5'-GACCACTACCAG-CAGAACAC-3') and 'eGFP-2-R' (5'-GAACTCCAG-CAGGACCATG-3'), and 200 nM probe 'eGFP-HEX' (5'-HEX-AGCACCCAGTCCGCCCTGAGCA-BHQ1-3') in a final volume of 25 µl.

### Outer membrane protein A genotyping PCR

Samples that were positive for *C. psittaci* in qPCR were subjected to an outer membrane protein A (*ompA*) genotyping PCR. Per sample, a reaction mix with a final volume of 50 µl containing 25 µl REDTaq ReadyMix (Merck KGaA), 200 nM of the primers 'ompA F (CTU)' (5'-ATGAAAAAACTCTTGAAATCGG-3') and 'ompA rev' (5'-TCCTTAGAATCTGAATTGAGC-3'), and 3 µl sample template with a DNA concentration of 25 ng/µl was prepared.<sup>31</sup> Cycling conditions were 10 minutes at 95°C, followed by 35 cycles of 95°C for 30 seconds, 49°C for 30 seconds, 72°C for 60 seconds and a final elongation at 72°C for seven minutes. PCR products were purified using the QIAquick PCR Purification Kit (Qiagen) according to the manufacturer's instructions. Purified amplicons were Sanger-sequenced by Microsynth. The obtained sequences were assembled and analysed using the Geneious Prime software version 2019.2.3 and compared against the National Center for Biotechnology Information database using the BLASTn tool (<https://blast.ncbi.nlm.nih.gov/>).

All primers and probes used in this study were obtained from Microsynth.

### RESULTS

*Chlamydiaceae* were detected in 0.9 per cent (95 per cent confidence interval: 0.3–2.6 per cent) of the birds (n=3) of three different species, namely in one of five (20 per cent, 3.6–62.5 per cent) fancy pigeons (*Columba livia domestica*), one of five (20 per cent, 3.6–62.5 per cent) common wood pigeons (*Columba palumbus*), and one of four (25 per cent, 4.6–69.9 per cent) Eurasian collared doves (*Streptopelia decaocto*), as shown in table 1. Five individual samples were positive for *Chlamydiaceae*: both choanal and cloacal swabs from the Eurasian collared dove and fancy pigeon, and a faecal swab from the common wood pigeon. Both choanal and cloacal swabs

**Table 1** Species, number of sample types and number of individual birds per total number of birds tested for *Chlamydiaceae* using real-time PCR in this study

Order	Family	Species name (Latin)	Species name (English)	Number of available and <i>Chlamydiaceae</i> -positive swab samples per anatomical site			<i>Chlamydiaceae</i> -positive birds (%; 95% CI)	Mean Ct value, <i>Chlamydiaceae</i> quantitative PCR	<i>Chlamydia</i> psittaci-positive birds (%)	Accession number of <i>ompA</i> study sequence
Anseriformes	Anatidae	<i>Anas platyrhynchos</i>	Mallard	Faeces	Choana	Cloaca	0/29 (0, 0–11.7)		n.d.	
		<i>Cygnus olor</i>	Mute swan	0/0	0/1	0/1	0/1 (0, 0–79.4)		n.d.	
		<i>Mergus merganser</i>	Common merganser	0/9	0/0	0/0	0/9 (0, 0–29.9)		n.d.	
Apodiformes	Apodidae	<i>Apus apus</i>	Common swift	0/15	0/7	0/7	0/20 (0, 0–16.1)		n.d.	
		<i>Tachymarptis melba</i>	Alpine swift	0/3	0/0	0/0	0/3 (0, 0–56.2)		n.d.	
Charadriiformes	Laridae	<i>Larus michahellis</i>	Yellow-legged gull	0/3	0/1	0/1	0/4 (0, 0–49.0)		n.d.	
Ciconiiformes	Ciconiidae	<i>Ciconia ciconia</i>	White stork	0/3	0/2	0/2	0/5 (0, 0–43.5)		n.d.	
Columbiformes	Columbidae	<i>Columba livia domestica</i>	Fancy pigeon	0/4	1/1	1/1	1/5 (20, 3.6–62.5)	Choana: 28.0 Cloaca: 29.3	1/1 (100)	MT450277
		<i>Columba livia domestica</i>	Feral pigeon	0/5	0/2	0/2	0/6 (0, 0–39.0)		n.d.	
		<i>Columba palumbus</i>	Common wood pigeon	1/4	0/1	0/1	1/5 (20, 3.6–62.5)	Faeces: 37.9	0/1 (0)	
Gruiformes	Rallidae	<i>Streptopelia decaocto</i>	Eurasian collared dove	0/2	1/2	1/2	1/4 (25, 4.6–69.9)	Choana: 36.0 Cloaca: 16.9	1/1 (100)	MT450278
		<i>Crex crex</i>	Corn crane	0/1	0/0	0/0	0/1 (0, 0–79.4)		n.d.	
		<i>Fulica atra</i>	Eurasian coot	0/2	0/1	0/1	0/3 (0, 0–56.2)		n.d.	
Passeriformes	Acrocephalidae	<i>Acrocephalus scirpaceus</i>	Eurasian reed warbler	0/0	0/1	0/1	0/1 (0, 0–79.4)		n.d.	
		<i>Emberiza citrinella</i>	Yellowhammer	0/1	0/1	0/1	0/1 (0, 0–79.4)		n.d.	
		<i>Carduelis carduelis</i>	Goldfinch	0/8	0/1	0/1	0/8 (0, 0–32.4)		n.d.	
Fringillidae	Fringillidae	<i>Chloris chloris</i>	European greenfinch	0/4	0/4	0/4	0/7 (0, 0–35.4)		n.d.	
		<i>Coccothraustes coccothraustes</i>	Hawfinch	0/1	0/1	0/1	0/2 (0, 0–65.8)		n.d.	
		<i>Fringilla coelebs</i>	Common chaffinch	0/4	0/1	0/1	0/4 (0, 0–49.0)		n.d.	
Hirundinidae	Hirundinidae	<i>Serinus serinus</i>	European serin	0/1	0/0	0/0	0/1 (0, 0–79.4)		n.d.	
		<i>Delichon urbicum</i>	Common house martin	0/2	0/0	0/0	0/2 (0, 0–65.8)		n.d.	
		<i>Hirundo rustica</i>	Barn swallow	0/4	0/2	0/2	0/6 (0, 0–39.0)		n.d.	
Motacillidae	Motacillidae	<i>Motacilla alba</i>	White wagtail	0/3	0/0	0/0	0/3 (0, 0–56.2)		n.d.	
		<i>Erithacus rubecula</i>	European robin	0/0	0/1	0/1	0/1 (0, 0–79.4)		n.d.	
		<i>Ficedula hypoleuca</i>	European pied flycatcher	0/1	0/0	0/0	0/1 (0, 0–79.4)		n.d.	
Muscicapidae	Muscicapidae	<i>Muscicapa striata</i>	Spotted flycatcher	0/3	0/0	0/0	0/3 (0, 0–56.2)		n.d.	
		<i>Phoenicurus ochruros</i>	Black redstart	0/6	0/2	0/2	0/8 (0, 0–32.4)		n.d.	
		<i>Cyanistes caeruleus</i>	Eurasian blue tit	0/18	0/7	0/7	0/25 (0, 0–13.3)		n.d.	
Paridae	Paridae	<i>Parus major</i>	Great tit	0/15	0/3	0/3	0/18 (0, 0–17.6)		n.d.	
		<i>Passer domesticus</i>	House sparrow	0/40	0/11	0/11	0/48 (0, 0–7.4)		n.d.	
		<i>Passer montanus</i>	Eurasian tree sparrow	0/5	0/1	0/1	0/6 (0, 0–39.0)		n.d.	

Continued

**Table 1** Continued

Order	Family	Species name (Latin)	Species name (English)	Number of available and <i>Chlamydiaceae</i> *-positive swab samples per anatomical site		<i>Chlamydiaceae</i> -positive birds (%; 95% CI)	Mean Ct value, <i>Chlamydiaceae</i> quantitative PCR	<i>Chlamydia psittaci</i> -positive birds (%)	Accession number of <i>ompA</i> study sequence
	Sittidae	<i>Sitta europaea</i>	Eurasian nuthatch	0/0	0/1	0/1 (0, 0–79.4)		n.d.	
	Sturnidae	<i>Sturnus vulgaris</i>	Common starling	0/4	0/2	0/6 (0, 0–39.0)		n.d.	
	Sylviidae	<i>Sylvia atricapilla</i>	Eurasian blackcap	0/7	0/1	0/7 (0, 0–35.4)		n.d.	
		<i>Sylvia borin</i>	Garden warbler	0/1	0/0	0/1 (0, 0–79.4)		n.d.	
	Turdidae	<i>Turdus merula</i>	Blackbird	0/50	0/22	0/21	0/67 (0, 0–5.4)	n.d.	
		<i>Turdus philomelos</i>	Song thrush	0/3	0/3	0/3	0/4 (0, 0–49.0)	n.d.	
		<i>Turdus pilaris</i>	Fieldfare	0/2	0/1	0/1	0/3 (0, 0–56.2)	n.d.	
		<i>Turdus viscivorus</i>	Mistle thrush	0/1	0/0	0/0	0/1 (0, 0–79.4)	n.d.	
Pelecaniformes	Ardeidae	<i>Ardea cinerea</i>	Grey heron	0/0	0/2	0/2 (0, 0–65.8)		n.d.	
Piciformes	Picidae	<i>Dendrocopos major</i>	Great spotted woodpecker	0/3	0/1	0/3 (0, 0–56.2)		n.d.	
		<i>Picus viridis</i>	European green woodpecker	0/3	0/0	0/1	0/4 (0, 0–49.0)	n.d.	
Total				1/267	2/91	2/91	3/339 (0.9, 0.3–2.6)	2/3 (66.7)	

\*All faecal swabs were obtained from living birds after defecation. Both choanal and cloacal swabs were available from 89 deceased birds. From one blackbird only a choanal swab was available. From one Eurasian blue tit a faecal and a choanal swab were available. Only a cloacal swab was obtained from one Eurasian blue tit and one European green woodpecker. All three swab types were obtained from 20 birds that died or were euthanased during treatment due to trauma or disease.

CI, confidence interval; Ct, cycle threshold; n.d., not determined; *ompA*, outer membrane protein A.



from the Eurasian collared dove and fancy pigeon were positive for *C. psittaci* by species-specific qPCR. *OmpA* genotyping classified the organism detected in the cloacal sample of the Eurasian collared dove as *C. psittaci* B. *C. psittaci* detected in the choanal swab of the fancy pigeon belonged to the *ompA* genotype E.

*C. psittaci* was not detected in the *Chlamydiaceae*-positive faecal swab of the common wood pigeon, and *ompA* genotyping was not successful due to low copy numbers (table 1). Thus, it was not possible to specify the detected *Chlamydiaceae* in this sample.

## DISCUSSION

The workers at the rehabilitation centre belong to the population at risk for zoonotic diseases transmitted by birds. Among all bird orders, Columbiformes and psittacine birds show the highest *Chlamydia* prevalence, ranging between 3.4 per cent and 50 per cent.<sup>22–23</sup> In this study, three of 20 (15 per cent, 5.2–36.2 per cent) of the Columbiformes were positive for *Chlamydiaceae*, whereof two were positive for *C. psittaci*. This is in accordance with the findings of other studies performed in Switzerland.<sup>20–21</sup> *C. psittaci* genotype B, which was found in one Eurasian collared dove, is the predominant genotype in the European pigeon population.<sup>22–23</sup> Genotype E, which was found in one fancy pigeon, infects a variety of avian species and is frequently found in pigeons worldwide.<sup>34–37</sup> Both genotypes are zoonotic, but human infection is mostly associated with genotype A causing a more severe course of disease.<sup>38–40</sup>

*Chlamydiaceae* were not detected in any other bird order included in this study. Partly, the low infection rate might be due to the selection of study samples. The majority of birds (n=246) were tested via faecal swabs only. Testing of faecal swabs has been shown to be a less sensitive method for detection of *Chlamydiaceae* compared with choanal swabs.<sup>41</sup> Furthermore, most birds were nestlings (n=163) or juveniles (n=126), which were previously shown to have lower *Chlamydiaceae* prevalence than adult birds.<sup>18</sup> However, these circumstances reflect the real conditions in the rehabilitation centre during the peak season. The findings in this study are in accordance with those of Zweifel and others,<sup>21</sup> reporting low infection rates of non-*C. psittaci*-*Chlamydiaceae* in songbirds (0.4 per cent) and waterfowl (4.3 per cent). In this study, excretion of *Chlamydiaceae* under stressful conditions in a bird rehabilitation centre was only present in pigeons. As pigeons mostly harbour *C. psittaci*, they may therefore pose a hazard to workers at rehabilitation centres. In order to minimise the risk of zoonotic chlamydial transmission, protective equipment (eg, gloves, masks) should always be used and appropriate hygiene measures (eg, washing hands) should be adhered to when handling wild birds.<sup>42</sup>

**Acknowledgements** The authors would like to thank the team of the rehabilitation centre of the Swiss Ornithological Institute, Sempach, for the collection of study samples, and Brigitte Sigrist from the NRGK, Vetsuisse Faculty, University of Zurich, for the excellent technical assistance.

**Contributors** SA, BV and NB designed the study. SS and PM performed the experiments. SS, HM and NW analysed the data. SS wrote the original draft. SA, BV, NB, HM, NW and PM reviewed and edited the original draft. All authors have read and agreed to the published version of the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Ethics approval** All animal housing and sampling were conducted in strict accordance with the Swiss law of animal welfare. None of the birds was killed for this study. The birds of which choanal and cloacal swabs were taken were euthanased due to incurable trauma or disease before sampling.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, an indication of whether changes were made, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Sandro Stalder <http://orcid.org/0000-0002-1077-2923>

Barbara Vogler <http://orcid.org/0000-0002-8330-5269>

## REFERENCES

- Sachse K, Kuehlewind S, Ruettinger A, et al. More than classical *Chlamydia psittaci* in urban pigeons. *Vet Microbiol* 2012;157:476–80.
- Vorimore F, Hsia R-C, Huot-Creasy H, et al. Isolation of a new *Chlamydia* species from the feral sacred IBIS (*Threskiornis aethiopicus*): *Chlamydia ibidis*. *PLoS One* 2013;8:e74823.
- Di Francesco A, Donati M, Laroucau K, et al. *Chlamydiae* in corvids. *Vet Rec* 2015;177:466.
- Guo W, Li J, Kaltenboeck B, et al. *Chlamydia gallinacea*, not *C. psittaci*, is the endemic chlamydial species in chicken (*Gallus gallus*). *Sci Rep* 2016;6:19638.
- Szymańska-Czerwińska M, Mitura A, Niemczuk K, et al. Dissemination and genetic diversity of chlamydial agents in Polish wildfowl: isolation and molecular characterisation of avian *Chlamydia abortus* strains. *PLoS One* 2017;12:e0174599.
- Laroucau K, Vorimore F, Aaziz R, et al. *Chlamydia buteonis*, a new *Chlamydia* species isolated from a red-shouldered hawk. *Syst Appl Microbiol* 2019;42:125997.
- Stokes HS, Martens JM, Chamings A, et al. Identification of *Chlamydia gallinacea* in a parrot and in free-range chickens in Australia. *Aust Vet J* 2019;97:398–400.
- Kaleta EF, Taday EMA. Avian host range of *Chlamydophila* spp. based on isolation, antigen detection and serology. *Avian Pathol* 2003;32:435–61.
- Shewen PE. Chlamydial infection in animals: a review. *Can Vet J* 1980;21:2–11.
- Vanrompay D, Ducatelle R, Haesebrouck F. *Chlamydia psittaci* infections: a review with emphasis on avian chlamydiosis. *Vet Microbiol* 1995;45:93–119.
- Bundesamt für Lebensmittelsicherheit und Veterinärwesen (BLV). Informationssystem Seuchenmeldungen InfoSM, 2020. Available: <https://www.infosm.blv.admin.ch> [Accessed 3 Apr 2020].
- Beeckman DSA, Vanrompay DCG. Zoonotic *Chlamydophila psittaci* infections from a clinical perspective. *Clin Microbiol Infect* 2009;15:11–17.
- Olsen B, Persson K, Broholm KA. PCR detection of *Chlamydia psittaci* in faecal samples from passerine birds in Sweden. *Epidemiol Infect* 1998;121:481–4.
- Vlahović K, Matica B, Bata I, et al. *Campylobacter*, *Salmonella* and *Chlamydia* in free-living birds of Croatia. *Eur J Wildl Res* 2004;50:127–32.
- Schwarzová K, Betáková T, Neméth J, et al. Detection of *Borrelia burgdorferi* sensu lato and *Chlamydophila psittaci* in throat and cloacal swabs from birds migrating through Slovakia. *Folia Microbiol* 2006;51:653–8.
- Sharples E, Baines SJ. Prevalence of *Chlamydophila psittaci*-positive cloacal PCR tests in wild avian casualties in the UK. *Vet Rec* 2009;164:16–17.

- 17 Krawiec M, Piasecki T, Wieliczko A. Prevalence of *Chlamydia psittaci* and other *Chlamydia* species in wild birds in Poland. *Vector Borne Zoonotic Dis* 2015;15:652–5.
- 18 Konicek C, Vondrážka P, Barták P, et al. DETECTION OF ZOONOTIC PATHOGENS IN WILD BIRDS IN THE CROSS-BORDER REGION AUSTRIA - CZECH REPUBLIC. *J Wildl Dis* 2016;52:850–61.
- 19 Geigenfeind I, Vanrompay D, Haag-Wackernagel D. Prevalence of *Chlamydia psittaci* in the feral pigeon population of Basel, Switzerland. *J Med Microbiol* 2012;61:261–5.
- 20 Mattmann P, Marti H, Borel N, et al. *Chlamydiaceae* in wild, feral and domestic pigeons in Switzerland and insight into population dynamics by *Chlamydia psittaci* multilocus sequence typing. *PLoS One* 2019;14:e0226088.
- 21 Zweifel D, Hoop R, Sachse K, et al. Prevalence of *Chlamydophila psittaci* in wild birds—potential risk for domestic poultry, pet birds, and public health? *Eur J Wildl Res* 2009;55:575–81.
- 22 Magnino S, Haag-Wackernagel D, Geigenfeind I, et al. Chlamydial infections in feral pigeons in Europe: review of data and focus on public health implications. *Vet Microbiol* 2009;135:54–67.
- 23 de Freitas Raso T, Seixas GHF, Guedes NMR, et al. *Chlamydophila psittaci* in free-living Blue-fronted Amazon parrots (*Amazona aestiva*) and Hyacinth macaws (*Anodorhynchus hyacinthinus*) in the Pantanal of Mato Grosso do Sul, Brazil. *Vet Microbiol* 2006;117:235–41.
- 24 Holzinger-Umlauf HA, Marschang RE, Gravendyck M, et al. Investigation on the frequency of *Chlamydia* sp. infections in tits (Paridae). *Avian Pathol* 1997;26:779–89.
- 25 Pennycott TW, Dagleish MP, Wood AM, et al. *Chlamydophila psittaci* in wild birds in the UK. *Vet Rec* 2009;164:157–8.
- 26 Ehrlich R, Slickers P, Goellner S, et al. Optimized DNA microarray assay allows detection and genotyping of single PCR-amplifiable target copies. *Mol Cell Probes* 2006;20:60–3.
- 27 Hoffmann B, Depner K, Schirrmeier H, et al. A universal heterologous internal control system for duplex real-time RT-PCR assays used in a detection system for pestiviruses. *J Virol Methods* 2006;136:200–9.
- 28 Blumer S, Greub G, Waldvogel A, et al. Waddlia, Parachlamydia and *Chlamydiaceae* in bovine abortion. *Vet Microbiol* 2011;152:385–93.
- 29 Pantchev A, Sting R, Bauerfeind R, et al. New real-time PCR tests for species-specific detection of *Chlamydophila psittaci* and *Chlamydophila abortus* from tissue samples. *Vet J* 2009;181:145–50.
- 30 Hoffmann B, Beer M, Schelp C, et al. Validation of a real-time RT-PCR assay for sensitive and specific detection of classical swine fever. *J Virol Methods* 2005;130:36–44.
- 31 Sachse K, Laroucau K, Hotzel H, et al. Genotyping of *Chlamydophila psittaci* using a new DNA microarray assay based on sequence analysis of *ompA* genes. *BMC Microbiol* 2008;8:63.
- 32 Dolz G, Solórzano-Morales A, Angelova L, et al. *Chlamydia psittaci* genotype B in a pigeon (*Columba livia*) inhabiting a public place in San José, Costa Rica. *Open Vet J* 2013;3:135–9.
- 33 Wang X, Zhang N-Z, Ma C-F, et al. Epidemiological investigation and genotype of *Chlamydia* exposure in pigeons in three provinces in northern China. *Vector Borne Zoonotic Dis* 2018;18:181–4.
- 34 Dickx V, Kalmar ID, Tavernier P, et al. Prevalence and genotype distribution of *Chlamydia psittaci* in feral Canada geese (*Branta canadensis*) in Belgium. *Vector Borne Zoonotic Dis* 2013;13:382–4.
- 35 Beckmann KM, Borel N, Pocknell AM, et al. Chlamydiosis in British garden birds (2005–2011): retrospective diagnosis and *Chlamydia psittaci* genotype determination. *Ecohealth* 2014;11:544–63.
- 36 Donati M, Laroucau K, Delogu M, et al. *Chlamydia psittaci* in Eurasian collared doves (*Streptopelia decaocto*) in Italy. *J Wildl Dis* 2015;51:214–7.
- 37 Jeong J, An I, Oem J-K, et al. Molecular prevalence and genotyping of *Chlamydia* spp. in wild birds from South Korea. *J Vet Med Sci* 2017;79:1204–9.
- 38 Rehn M, Ringberg H, Runeheger A, et al. Unusual increase of psittacosis in southern Sweden linked to wild bird exposure, January to April 2013. *Euro Surveill* 2013;18:20478.
- 39 Wallensten A, Fredlund H, Runeheger A. Multiple human-to-human transmission from a severe case of psittacosis, Sweden, January–February 2013. *Euro Surveill* 2014;19:20937.
- 40 Cadario ME, Frutos MC, Arias MB, et al. Epidemiological and molecular characteristics of *Chlamydia psittaci* from 8 human cases of psittacosis and 4 related birds in Argentina. *Rev Argent Microbiol* 2017;49:323–7.
- 41 Andersen AA. Comparison of pharyngeal, fecal, and cloacal samples for the isolation of *Chlamydia psittaci* from experimentally infected cockatiels and turkeys. *J Vet Diagn Invest* 1996;8:448–50.
- 42 Tolba HMN, Abou Elez RMM, Elsohaby I. Risk factors associated with *Chlamydia psittaci* infections in psittacine birds and bird handlers. *J Appl Microbiol* 2019;126:402–10.